

Title:

Full range of motion induces greater muscle damage than partial range of motion in elbow flexion exercise with free weights

Running title:

Range of motion affects exercise-induced muscle damage

Laboratory where the research was conducted:

Exercise Research Laboratory, Federal University of Rio Grande do Sul, Brazil

Authors:

Bruno Manfredini Baroni¹⁻², Marcelo Gava Pompermayer¹, Anelize Cini¹, Amanda Storti Peruzzolo¹, Régis Radaelli¹, Clarissa Müller Brusco¹, Ronei Silveira Pinto¹

Authors' affiliations:

¹ Strength Training Research Group, Exercise Research Laboratory, Federal University of Rio Grande do Sul, Brazil

² Graduate Program in Rehabilitation Sciences, Federal University of Health Sciences of Porto Alegre, Brazil

Corresponding Author:

Bruno Manfredini Baroni

Federal University of Health Sciences of Porto Alegre

Sarmento Leite St., 245

Porto Alegre, Rio Grande do Sul, Brazil

Postal code: 90050-170

Phone: +55 (51) 3303-8876

E-mail: bmbaroni@yahoo.com.br

ABSTRACT

Load and range of motion (ROM) applied in resistance training (RT) affect the muscle damage magnitude and the recovery time-course. Since exercises performed with partial ROM allow a higher load compared to those with full ROM, this study investigated the acute effect of a traditional RT exercise using full ROM or partial ROM on muscle damage markers. Fourteen healthy men performed four sets of 10 concentric-eccentric repetitions of unilateral elbow flexion at the Scott bench. Arms were randomly assigned to partial ROM (50° to 100°) and full ROM (0° to 130°) conditions, and load was determined as 80% of one repetition maximum in full and partial ROM tests. Muscle damage markers were assessed pre-exercise, and immediately, 24h, 48h and 72h after exercise. Primary outcomes were peak torque, muscle soreness during palpation and elbow extension, arm circumference, and joint ROM. The load lifted in partial ROM condition (1RM=19.1±3.0kg) was 40±18% higher compared to full ROM (1RM=13.7±2.2kg). 72h after exercise, the full ROM condition led to significant higher soreness sensation during elbow extension (1.3-4.1cm vs. 1.0-1.9cm) and smaller ROM values (97.5-106.1° vs. 103.6-115.7°). Peak torque, soreness from palpation and arm circumference were statistically similar between conditions, although mean values in all time points of these outcomes have suggested a more expressive muscle damage for the full ROM condition. In conclusion, elbow flexion exercise with full ROM seems to induce greater muscle damage than partial ROM exercises, even though higher absolute load was achieved with partial ROM.

Keywords: Strength training, Exercise-induced muscle damage, Scott bench.

INTRODUCTION

Resistance training (RT) is often reported as the most effective method for long-term increases in strength and muscle mass (31). However, acute effects of RT should be considered for prescription and periodization, since individuals participating in RT experience exercise-induced muscle damage, which impairs muscle function and limits performance for some days following exercise (7). Therefore, sufficient recovery time should be implemented between training sessions, which could be achieved in a gym environment through assessment of indirect markers of muscle damage, such as muscle strength capacity, delayed-onset muscle soreness (DOMS), joint range of motion (ROM) and limb circumference (CIR) (7, 34).

Mechanical factors appear to initiate the injury process, while a cascade of metabolic events and inflammation increase muscle damage in the following days after exercise (24). Exercise intensity (i.e. load) and volume (i.e. number of sets and repetitions) are the most commonly manipulated variables in RT programs, which affect mechanical stress over the muscle and, consequently, the magnitude of exercise-induced muscle damage (5, 20). However, other intervenient factors on mechanical stress applied over the muscle tissue should be considered, such as: muscle group (3), type of exercise (30), type of muscle action (12) and movement velocity (2). Additionally, evidence suggests that joint ROM employed during resistance exercise may also affect the muscle damage response (11, 17, 23).

Previous studies (11, 17, 23) showed greater muscle damage (or a trend to) in elbow flexor muscles exercised at long compared to short muscle lengths. However, these studies have exclusively used maximal eccentric contractions, performed in isokinetic conditions, and with a fixed ROM starting at short or long muscle lengths. Importantly, these conditions are dissimilar from those observed during a traditional RT program (i.e. using free weights or gym machines). Exercises with full ROM have been recommended in traditional RT, despite a partial ROM (when long and short muscle lengths are avoided) allows an exercise to be accomplished with a higher load (13, 14, 25). In fact, weightlifters and RT practitioners commonly perform exercises with partial ROM in order to displace superior amounts of load or even to complete a target

number of repetitions during a set. However, there is a lack in the literature regarding the acute effects of ROM on muscle damage induced by traditional RT exercises.

In summary, it remains unknown whether the higher load lifted in the partial ROM condition or the larger angular amplitude during the full ROM exercise is the preponderant factor for muscle damage magnitude and recovery time-course in traditional RT exercise. Therefore, the purpose of this study was to investigate the acute effect of a traditional RT exercise using full ROM or partial ROM on muscle damage markers of healthy young men. Taking into consideration previous findings that full ROM seemed to promote higher strength gains compared to partial ROM after a systematized RT program (13, 25), and assuming a close relationship between muscle damage generated in RT sessions and muscular adaptation (29), we hypothesized that muscle damage would be more expressive at the full ROM condition.

METHODS

Experimental approach to the problem

After recruitment, the subjects visited the laboratory for data collection on seven occasions (Figure 1). In the first visit, they were familiarized to the procedures and tests. In the second visit, one repetition maximum (1RM) test was performed; one arm performed the elbow flexion 1RM test with a partial ROM, while the contralateral arm performed the elbow flexion 1RM test with a full ROM, according to the randomization protocol. Although evidence suggests similar responses of preferred and non-preferred arms to a muscle damage protocol (19), an equal number of volunteers performed each experimental condition at the preferred side. After 48h, both arms repeated the 1RM test for reliability purposes. After at least 72h, subjects returned to the laboratory to perform the respective RT protocol determined for each arm. The indirect markers of damage were assessed before and immediately after exercise, as well as 24h, 48h and 72h after exercise in both arms. The study was conducted according to the declaration of Helsinki and all procedures were approved by the local Research Ethics Committee.

<< Figure 1 near here >>

Participants

Fourteen untrained male undergraduate subjects, aged between 18 and 25 years, volunteered for the study. As a part of the inclusion criteria, subjects were not to be involved in any systematized training programs for at least three months before they started the study. All subjects were informed about the purpose, procedures, benefits and risks that might result from this study, and agreed to participate through a signed informed consent. Physical characteristics (mean \pm SD) for age, weight and height were 21.1 \pm 1.9 years, 77.1 \pm 13.6 kg and 178.9 \pm 5.6 cm, respectively. Participants were instructed to perform no vigorous physical activity during the study and not to take any medication, dietary supplement or alcoholic beverages.

One repetition maximum test

To determine the 1RM load, subjects visited the laboratory two times with at least 48h between each session. The same procedures were maintained on both occasions. Subjects warmed-up with a light to moderate load for 10 repetitions of unilateral elbow flexion using a dumbbell resistance on Scott bench (preacher curl, Sculptor, Porto Alegre, RS, Brazil). According to the randomization protocol, one arm was tested at partial ROM (50° to 100°) and the other at full ROM (0° to 130°), considering 0° as full elbow extension. To control ROM, two metallic bars were used to regulate dumbbell displacement (25). Adjustments to determine 1RM load were made through trial-and-error, as previously described (27). No more than five attempts with a 3-min rest interval between trials were necessary to determine 1RM. A metronome controlled the cadence: 1s for each phase (concentric and eccentric) for the partial ROM condition, and 2s for each phase for the full ROM condition. The 1RM test was ceased if the subject was unable to move the dumbbell in the target cadence.

Resistance training session

The RT protocol consisted of four sets of 10 concentric-eccentric repetitions on unilateral elbow flexion using a dumbbell resistance equal to 80% of full or partial ROM 1RM at the same Scott bench used for 1RM tests. Cadence was controlled through a metronome: 1s for each phase for the partial

ROM condition; and 2s for each phase for the full ROM condition. A 2-min rest interval was used between sets. According to previous randomization and 1RM tests, one arm performed the elbow flexion exercise at partial ROM (50°-100°) while the contralateral arm performed the exercise at a full ROM (0°-130°). Importantly, some volunteers were not able to perform 130° of ROM due to limited elbow flexion by the muscle mass and/or to limited elbow extension by the reduced stretching or fatigue and swelling during exercise; however, all participants in full ROM condition performed at least two fold more range of motion than the partial ROM condition.

Estimated work

Trigonometric functions were used to calculate the resistance torque for each joint position (resolution = 1°) in the exercise in full and partial ROM conditions. The mean value of subjects' forearm length (30 cm), the forearm position in relation to the ground, the mean load supported in 1RM for each experimental condition and the gravity effect were considered for resistance torque calculations. Movement acceleration effects were not considered in the calculations. Torque-angle curves were plotted and the areas under curves were considered the estimated work performed in full ROM and partial ROM conditions.

Peak torque

Elbow flexion peak torque (PT) was assessed using a Cybex NORM dynamometer (Ronkonkoma, NY, USA) attached to a Scott bench, as described elsewhere (8). Briefly, subjects were positioned in the Scott bench and the dynamometer rotation center was aligned to the elbow rotation center. The test consisted of three maximum isometric voluntary contractions of the elbow flexors. Elbow was kept in at 90°, each contraction lasted 5 seconds and a 2-min rest interval was maintained between tests. Subjects were verbally encouraged to maximal effort during all tests. The greatest PT among three trials was used for further analysis.

Muscle soreness

Muscle soreness was evaluated using a visual analog scale, which consisted of a line with 100 mm without any numbers or markers. The left end of the line denoted “no pain” while the right end represented “extremely painful”. Subjects were instructed to draw a mark on the line according to their sensation during two situations: elbow extension movement; and palpation performed by a researcher at the middle point of muscle belly of the biceps brachii. Soreness was quantified by measuring the distance between the initial point line (“no pain”) and the point marked by the subject (1).

Arm circumference

Subjects remained standing with the arms relaxed beside their body for circumference (CIR) measurement. The CIR of the upper arm was determined by placing a tape measure at 60% of the distance between lateral epicondyle and acromion. The CIR measurement sites were marked with a semi permanent ink pen to ensure reliability in subsequent assessments (3).

Range of motion

A plastic goniometer (Carci, SP, Brazil) was used to assess ROM. Subjects remained at standing position with the arms relaxed for goniometer positioning, and the goniometer was placed on the elbow joint at the lateral epicondyle of the humerus. Then, subjects were asked to maximally flex the elbow. ROM consisted of the total angular amplitude starting from elbow in the relaxed position until maximal voluntary elbow flexion(3).

Statistical analysis

A paired sample t test was used to compare 1RM load between conditions. A Two-Way ANOVA (condition x time) was used to compare PT, soreness, ROM and CIR. *A priori* sphericity was tested through Mauchly’s test, and Greenhouse-Geisser correction was used when necessary. Statistical significance was set at an alpha level of 0.05. Results are expressed as mean \pm standard deviation. All analyses were performed at SPSS version 20.0.

RESULTS

All subjects were able to complete the number of sets and repetitions determined in both experimental conditions at the RT session. Significantly higher loads ($40\pm 18\%$; $p < 0.01$) were lifted in the partial ROM condition ($1RM = 19.07 \pm 3.02$ kg) compared to the full ROM condition ($1RM = 13.71 \pm 2.16$ kg). Considering these mean load values for each exercise condition, the full ROM condition led to a larger estimated work compared to the partial ROM condition (Figure 2).

<< Figure 2 near here >>

PT presented no time-condition interaction ($p = 0.204$), and just a significant time effect was observed ($p < 0.001$). PT decreased over all time points and remained significantly lower than baseline up to 72h after exercise ($p < 0.001$). The lowest PT values occurred at 24h for both conditions. PT fall ranged between 23-29% at the full ROM condition and 9-22% at the partial ROM condition throughout the study (Figure 3).

<< Figure 3 near here >>

Muscle soreness at elbow extension presented a significant time-condition interaction ($p = 0.031$). This soreness test showed significantly higher values for full ROM compared to partial ROM at 48h ($p = 0.048$) and 72h ($p = 0.012$). In addition, soreness at the full ROM arm remained higher than baseline values until 72h ($p < 0.001$), while no significant changes occurred at the partial ROM arm along time ($p > 0.05$ for all time point comparisons). Peak soreness levels were found at 24h for partial ROM condition and at 48h for full ROM condition. Mean soreness values at the full ROM arm were 31-248% higher than the partial ROM arm between 24h and 72h after exercise (Figure 4 – left graph).

Muscle soreness during palpation had no time-condition interaction ($p = 0.148$), and just a significant time effect was observed ($p < 0.001$). A significant increase in all time points compared to baseline ($p < 0.01$) was found, except for immediately after exercise ($p = 0.670$). Peak values occurred at 24h post-exercise in both conditions. Mean soreness values at the full ROM arm were 76-129% higher than the partial ROM condition between 24h and 72h after exercise (Figure 4 – right graph).

<< Figure 4 near here >>

CIR also had no time-condition interaction ($p=0.279$), but a significant time effect ($p<0.001$). CIR demonstrated significant increment in all time points compared to baseline ($p<0.05$), except for 48h post-exercise ($p=0.322$). The highest values were seen immediately after exercise for both conditions, and CIR increased 1-3% throughout the data collection (Figure 5).

<< Figure 5 near here >>

ROM presented a significant time-condition interaction ($p=0.029$). This outcome was significantly lower in the full ROM condition compared to the partial ROM at 48h ($p=0.007$) and 72h ($p=0.029$). Values in the full ROM condition were significantly reduced compared to baseline up to 72h ($p<0.01$), while partial ROM showed a significant decrease only immediately after exercise ($p=0.003$). The lowest ROM values were found immediately after exercise for both conditions. ROM decreases between 3-13% and 12-19% at partial and full ROM conditions, respectively (Figure 6).

<< Figure 6 near here >>

DISCUSSION

To the best of our knowledge, this is the first study designed to compare muscle damage in elbow flexors following a traditional RT session with different angular amplitudes (full ROM and partial ROM) and, consequently, different loads. In summary, our findings suggest that ROM during RT of elbow flexors seems to be as or even more important than load itself for muscle damage magnitude, since the higher loads in partial ROM condition did not lead to greater levels of muscle damage compared to full ROM condition in any outcome.

Subjects' characteristics, such as gender (10), age (28) and conditioning status (18), as well as the exercised muscle group (lower vs. upper limb) (3), type of exercise (e.g., single- vs. multi-joint exercises) (30), type of muscle action (e.g., concentric vs. eccentric)(12), contraction velocity (2), exercise intensity/load (20) and number of repetitions (4) are variables that notably affect the muscle damage magnitude. In our study, elbow flexor muscles from each subject performed a similar exercise (Scott bench), with a controlled angular velocity (about 50-65°/s) and a standard training volume (four sets of 10 repetitions). As expected, participants endured a 40% higher load in the partial

ROM as compared to full ROM, further supporting findings involving the elbow flexion (25) and bench press (13, 14) exercises. Nonetheless, Figure 2 of our study makes it clear that the superior load in the partial ROM condition is insufficient to match the amount of work performed at the full ROM exercise. This is the practical condition that we had intention to test, i.e. the sum of both intervenient factors (joint ROM and exercise load), as commonly experienced by RT practitioners.

Nosaka and Sakamoto (23) compared isokinetic eccentric exercise bouts performed at different starting joint angles but with a similar ROM (80°), and found that longer muscle lengths induced greater muscle damage responses. While a similar peak torque among conditions was achieved, the arm performed a higher total amount of work while at a shorter muscle length (23). Therefore, although the higher estimated work performed by our volunteers in the full ROM compared to the partial ROM condition, the post-exercise response observed here seems to be mainly attributed to the longer muscle length achieved with full ROM. This relationship between fiber length and muscle damage seems very reasonable, and has been supported by experimental studies since the “popping-sarcomere hypothesis”, proposed by Morgan (15), according to which the sarcomeres are more susceptible to injury when exercised on the descending limb of the force-length relationship.

Previous studies focused in the ROM effect on muscle damage were restricted to isokinetic eccentric contractions (11, 17, 23), which is not the usual approach for traditional RT. Therefore, until now, it was unknown whether the higher loads used in the partial ROM condition would be sufficient to compensate the larger muscle damage induced by exercises performed at longer muscle lengths in the full ROM condition. According to our findings, full ROM exercise seems to promote higher muscle damage levels, even though with a smaller absolute load compared to the partial ROM exercise. However, this conclusion is not statistically supported by all outcomes of this study (i.e. PT, DOMS at palpation and CIR). This specific magnitude and time-course response of each muscle damage marker has already been discussed elsewhere (7), and that is the reason why studies involving exercise-induced muscle damage usually employ three or more outcomes in their experimental approaches.

In fact, isometric PT is referred in the literature as the most reliable marker of muscle damage (34), since maximal strength capacity is commonly used as representative of muscle function. We could not observe significant differences in PT values between our two experimental conditions, suggesting a similar level of exercise-induced muscle damage. However, it is interesting to note that the full ROM arm displayed strength falls around 23-29% throughout post-exercise days, while the partial ROM arm presented falls ranging between 9-22% in the same period. So, we believe the large data dispersion impaired our PT results, and it is possible that a larger sample size would enable the rejection of the null hypothesis (16).

DOMS is one of the most used markers of muscle damage (34), although it does not fully reflect the magnitude of microscopic injury to muscle cells (21). Since this is a completely subjective outcome and characterized by high dispersion between subjects (1), we chose two ways to assess muscle soreness: elbow extension and palpation. The results reported here are in agreement with those of Nosaka and Sakamoto (23), who reported that elbow extension triggers greater muscle soreness than local palpation, and soreness response is more pronounced at muscles exercised in longer lengths. Although our statistical approach indicated no time-condition interaction for DOMS during palpation, the full ROM arm presented mean values 76-129% higher than the contralateral arm. It should be highlighted that a 30% difference in pain scores is considered enough to be a clinically important response (35), thereby further supporting the higher DOMS induced by the full ROM condition in our study.

DOMS during elbow extension may be at least partially explained in our study by the more expressive ROM reduction observed following the full ROM condition. Increased muscle stiffness following damaging exercises is often reported (12), and it imposes difficulty for subjects in extending the elbow joint, which may account for painful sensation. At the same time, both DOMS and ROM may be related to muscle swelling, measured here through arm CIR. However, a more precise method to assess muscle swelling may be necessary to draw conclusions about this muscle damage marker, such as ultrasound images to assess muscle thickness and echo intensity (27).

Independent of the responsible mechanism for increased DOMS and reduced ROM, one factor should be taken into account by coaches and trainers:

DOMS and ROM are related to subjects' functional capacity. An uncomfortable situation promoted by a painful muscle group may lead to biomechanical changes in daily and sport tasks (6), potentially leading to decreased performance and increased injury risk. Similar consequences may be attributed to a reduced ROM capacity (33). In a simple example based in our own findings, muscles subjected to the full ROM condition would be unable to complete a subsequent bout using the same exercise with full ROM up to 72h after exercise, since subjects were not able to reach the last degrees of elbow extension without a certain level of discomfort.

Athletes and other RT practitioners frequently execute a training session without full muscle recovery from the previous one. Although muscle damage does not appear to be exacerbated following a RT session performed with insufficient recovery (5), it is likely that such a chronic model would lead to increased injury risk (6). In order to prevent the onset of maladaptation, it is important for athletes and active populations, that coaches and trainers track their recovery status for proper prescription. In this way, the inexpensive and practical assessments used in this study (i.e., DOMS, ROM and CIR) are available for these professionals.

RT experts usually recommend complete ROM during exercise. Since the most common goal of RT is to enhance muscle strength capacity, and considering that muscle damage level is usually considered a determinant factor for muscle hypertrophy and consequent strength improvement (29), our findings support this recommendation. However, an appropriate periodization using this condition should encompass sparse training sessions of the same muscle group into a week for a sufficient time for muscle recovery. On the other hand, a partial ROM bout appears to confer a protective effect on a subsequent full ROM bout (22); so, partial ROM exercises could be used before those with full ROM as a training progression in beginners or individuals who should avoid high levels of muscle damage. Furthermore, it is suggested that long-term interventions with partial ROM exercise induces specific strength gains near to the joint angles achieved in training (9), while the exercise executed at long muscle length led an increase in serial sarcomeres (sarcomerogenesis) (26), providing higher shortening velocity to muscle fibers (32). Therefore, specificity

of the subject's activity and goals should also be taking into consideration for appropriated prescription.

A previous study from our research group investigated the effects of a 10-week RT program with partial and full ROM exercises on strength and muscle mass of elbow flexors (25), and demonstrated higher effect sizes for the full ROM condition. If we simply take these previous findings (25) with those of the current study, it could be hypothesized that the mechanism responsible for the more pronounced chronic adaptation in full ROM condition was the greater muscle damage. However, since these studies were conducted separately and we did not control other intervenient factors on muscle adaptation to RT, such as protein synthesis and hormonal response, it would be presumptuous to assume a causal relationship. Therefore, we encourage more studies investigating the mechanism for the responses in subjects trained with exercises encompassing full and partial ROM.

In summary, we concluded that elbow flexion exercise performed with full ROM induces greater muscle damage than partial ROM exercise, even though the latter allows a higher absolute load compared to the first one. These results were first demonstrated by our study, and should be considered to optimize RT prescription and periodization, respecting the magnitude and time-course of muscle damage and recovery following exercises with different ROM.

PRACTICAL APPLICATIONS

Traditional RT using different ROM led to dissimilar muscle damage and recovery. Therefore, prescription based on different ROM should consider specific goals (i.e. increase in strength at specific angles or at entire ROM), training schedule (i.e. number of weekly sessions and interval between sessions) and contraindication to muscle damage (e.g. individuals with increased risk for rhabdomyolysis). In addition, following the RT model tested here, coaches should consider a larger recovery period for muscle regeneration in the full ROM compared to the partial ROM bout. The assessment of common indirect markers of muscle damage (i.e. force, ROM, DOMS and CIR) for tracking recovery status could be helpful for coaches to take reasonable decisions.

REFERENCES

1. Baroni BM, Leal Junior EC, De Marchi T, Lopes AL, Salvador M, and Vaz MA. Low level laser therapy before eccentric exercise reduces muscle damage markers in humans. *Eur J Appl Physiol* 110: 789-796, 2010.
2. Chapman D, Newton M, Sacco P, and Nosaka K. Greater muscle damage induced by fast versus slow velocity eccentric exercise. *Int J Sports Med* 27: 591-598, 2006.
3. Chen TC, Lin KY, Chen HL, Lin MJ, and Nosaka K. Comparison in eccentric exercise-induced muscle damage among four limb muscles. *Eur J Appl Physiol* 111: 211-223, 2011.
4. Chen TC and Nosaka K. Effects of number of eccentric muscle actions on first and second bouts of eccentric exercise of the elbow flexors. *J Sci Med Sport* 9: 57-66, 2006.
5. Chen TC and Nosaka K. Responses of elbow flexors to two strenuous eccentric exercise bouts separated by three days. *J Strength Cond Res* 20: 108-116, 2006.
6. Cheung K, Hume P, and Maxwell L. Delayed onset muscle soreness : treatment strategies and performance factors. *Sports Med* 33: 145-164, 2003.
7. Clarkson PM and Hubal MJ. Exercise-induced muscle damage in humans. *Am J Phys Med Rehabil* 81: S52-69, 2002.
8. Gentil P, Soares SR, Pereira MC, da Cunha RR, Martorelli SS, Martorelli AS, and Bottaro M. Effect of adding single-joint exercises to a multi-joint exercise resistance-training program on strength and hypertrophy in untrained subjects. *Appl Physiol Nutr Metab* 38: 341-344, 2013.
9. Graves JE, Pollock ML, Jones AE, Colvin AB, and Leggett SH. Specificity of limited range of motion variable resistance training. *Med Sci Sports Exerc* 21: 84-89, 1989.
10. Heavens KR, Szivak TK, Hooper DR, Dunn-Lewis C, Comstock BA, Flanagan SD, Looney DP, Kupchak BR, Maresh CM, Volek JS, and Kraemer WJ. The effects of high intensity short rest resistance exercise on muscle damage markers in men and women. *J Strength Cond Res* 28: 1041-1049, 2014.
11. Jones DA, Newham DJ, and Torgan C. Mechanical influences on long-lasting human muscle fatigue and delayed-onset pain. *J Physiol* 412: 415-427, 1989.
12. Lavender AP and Nosaka K. Changes in fluctuation of isometric force following eccentric and concentric exercise of the elbow flexors. *Eur J Appl Physiol* 96: 235-240, 2006.
13. Massey CD, Vincent J, Maneval M, and Johnson JT. Influence of range of motion in resistance training in women: early phase adaptations. *J Strength Cond Res* 19: 409-411, 2005.

14. Massey CD, Vincent J, Maneval M, Moore M, and Johnson JT. An analysis of full range of motion vs. partial range of motion training in the development of strength in untrained men. *J Strength Cond Res* 18: 518-521, 2004.
15. Morgan DL. New insights into the behavior of muscle during active lengthening. *Biophys J* 57: 209-221, 1990.
16. Nakagawa S and Cuthill IC. Effect size, confidence interval and statistical significance: a practical guide for biologists. *Biol Rev Camb Philos Soc* 82: 591-605, 2007.
17. Newham DJ, Jones DA, Ghosh G, and Aurora P. Muscle fatigue and pain after eccentric contractions at long and short length. *Clin Sci (Lond)* 74: 553-557, 1988.
18. Newton MJ, Morgan GT, Sacco P, Chapman DW, and Nosaka K. Comparison of responses to strenuous eccentric exercise of the elbow flexors between resistance-trained and untrained men. *J Strength Cond Res* 22: 597-607, 2008.
19. Newton MJ, Sacco P, Chapman D, and Nosaka K. Do dominant and non-dominant arms respond similarly to maximal eccentric exercise of the elbow flexors? *J Sci Med Sport* 16: 166-171, 2013.
20. Nosaka K and Newton M. Difference in the magnitude of muscle damage between maximal and submaximal eccentric loading. *J Strength Cond Res* 16: 202-208, 2002.
21. Nosaka K, Newton M, and Sacco P. Delayed-onset muscle soreness does not reflect the magnitude of eccentric exercise-induced muscle damage. *Scand J Med Sci Sports* 12: 337-346, 2002.
22. Nosaka K, Newton M, Sacco P, Chapman D, and Lavender A. Partial protection against muscle damage by eccentric actions at short muscle lengths. *Med Sci Sports Exerc* 37: 746-753, 2005.
23. Nosaka K and Sakamoto K. Effect of elbow joint angle on the magnitude of muscle damage to the elbow flexors. *Med Sci Sports Exerc* 33: 22-29, 2001.
24. Paulsen G, Mikkelsen UR, Raastad T, and Peake JM. Leucocytes, cytokines and satellite cells: what role do they play in muscle damage and regeneration following eccentric exercise? *Exerc Immunol Rev* 18: 42-97, 2012.
25. Pinto RS, Gomes N, Radaelli R, Botton CE, Brown LE, and Bottaro M. Effect of range of motion on muscle strength and thickness. *J Strength Cond Res* 26: 2140-2145, 2012.
26. Proske U and Morgan DL. Muscle damage from eccentric exercise: mechanism, mechanical signs, adaptation and clinical applications. *J Physiol* 537: 333-345, 2001.
27. Radaelli R, Bottaro M, Wilhelm EN, Wagner DR, and Pinto RS. Time course of strength and echo intensity recovery after resistance exercise in women. *J Strength Cond Res* 26: 2577-2584, 2012.
28. Roth SM, Martel GF, Ivey FM, Lemmer JT, Metter EJ, Hurley BF, and Rogers MA. High-volume, heavy-resistance strength training and muscle damage in young and older women. *J Appl Physiol (1985)* 88: 1112-1118, 2000.

29. Schoenfeld BJ. The mechanisms of muscle hypertrophy and their application to resistance training. *J Strength Cond Res* 24: 2857-2872, 2010.
30. Soares S, Ferreira-Junior JB, Pereira MC, Cleto VA, Castanheira RP, Cadore EL, Brown LE, Gentil P, Bembem MG, and Bottaro M. Dissociated Time Course of Muscle Damage Recovery Between Single- and Multi-Joint Exercises in Highly Resistance-Trained Men. *J Strength Cond Res* 29: 2594-2599, 2015.
31. Stone M, Plisk S, and Collins D. Training principles: evaluation of modes and methods of resistance training--a coaching perspective. *Sports Biomech* 1: 79-103, 2002.
32. Thom JM, Morse CI, Birch KM, and Narici MV. Influence of muscle architecture on the torque and power-velocity characteristics of young and elderly men. *Eur J Appl Physiol* 100: 613-619, 2007.
33. Verrall GM, Slavotinek JP, Barnes PG, Esterman A, Oakeshott RD, and Spriggins AJ. Hip joint range of motion restriction precedes athletic chronic groin injury. *J Sci Med Sport* 10: 463-466, 2007.
34. Warren GL, Lowe DA, and Armstrong RB. Measurement tools used in the study of eccentric contraction-induced injury. *Sports Med* 27: 43-59, 1999.
35. Yelland MJ and Schluter PJ. Defining worthwhile and desired responses to treatment of chronic low back pain. *Pain Med* 7: 38-45, 2006.

ACKNOWLEDGMENTS

The authors would like to thank Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Fundação de Amparo à Pesquisa do Estado do Rio Grande do Sul (FAPERGS) and Conselho Nacional de Pesquisa (CNPq-Brazil) for financial support.

FIGURE LEGENDS

Figure 1. Simplified experimental design.

Figure 2. Resistance torque (calculated through mean values of forearm length and load) along the range of motion and estimated work (area under curve) performed in each condition.

Figure 3. Peak torque (PT) throughout testing sessions for both conditions.


significantly lower than pre-exercise for both conditions.


Figure 4. Muscle soreness during elbow extension (left) and palpation (right) throughout testing sessions for both conditions. \$ significantly higher than pre-exercise for full ROM; # significantly higher than pre-exercise for both conditions. * full ROM significantly higher than partial ROM.

Figure 5. Arm circumference (CIR) throughout testing sessions for both conditions. # significantly higher than pre-exercise for both conditions.


Figure 6. Range of motion (ROM) throughout testing sessions for both conditions. \$ significantly lower than pre-exercise for full ROM; & significantly lower than pre-exercise for partial ROM; * full ROM significantly lower than partial ROM.

ACCEPTED








ACCEPTED




ACCEPTED



ACCEPTED



ACCEPTED



ACCEPTED